

LASER DEFINED VISION

POSTOPERATIVE INSTRUCTIONS – THE DAY OF SURGERY

EYE DROPS • • • • •



ZYMAXID

Antibiotic to protect your eye against infection.



DUREZOL

Anti-inflammatory to help reduce swelling.
SHAKE BOTTLE BEFORE USE.



REFRESH PLUS

Lubricating drops for comfort.

THE DAY OF SURGERY • • • • •

Immediately go home and remove the shields to instill the first set of drops at 1 minute intervals:

- Durezol* (pink top) x 1
- Zymaxid 0.5%* (tan top) x 1

Replace the shields and take a nap or relax with your eyes closed for at least 2-3 hours.

Whenever you awaken, repeat the above drops 3 times per day.

You also have **Refresh** tears in the bag to use as needed for irritation or dryness.

POST OPERATIVE DAY 1 • • • • •

The next morning, you may remove the shields and return to the office for follow-up as directed. Use the drops as directed the morning prior to seeing Dr. Stonecipher as well.

FOLLOWING POST-OP DAY 1 • • • • •

The suggested regimen is:

- Durezol* (pink top)
- Zymaxid 0.5%* (tan top)
- 3x/day for at least 7 full days; then stop.

Refresh Tears - Use 1 drop in the surgical eye(s)
4x/day for at least 2 months.

To avoid washing out the previously applied drops, **wait approximately 1 minute between drops.**
***If you receive generic eye drops, the bottles may look different than the pictures shown above.**

LASER DEFINED VISION

POSTOPERATIVE PRK INSTRUCTIONS – THE DAY OF SURGERY

EYE DROPS • • • • •



ZYMAXID

Antibiotic to protect your eye against infection.



ACUVAIL

Anti-inflammatory to help with pain.



FML

Anti-inflammatory to help reduce swelling.



REFRESH PLUS

Lubricating drops for comfort.

USE ONLY FOR 48 HOURS. SHAKE BOTTLE BEFORE USE.

THE DAY OF SURGERY • • • • •

Immediately go home and instill the first set of drops at 1 minute intervals:

- FML 0.1%** (purple label) x 1
- Zymaxid 0.5%** (tan top) x 1

Whenever you awaken, repeat the above drops at roughly 2-3 hour intervals while awake. You also have **Refresh** tears in the bag to use as needed for irritation or dryness.

POST OPERATIVE DAY 1 • • • • •

The next morning, you will return to the office for follow-up as directed. Use the drops as directed the morning prior to seeing Dr. Stonecipher as well.

FOLLOWING POST-OP DAY 1 • • • • •

The suggested regimen is:

- Zymaxid** (Antibiotic) – 1 drop in each operative eye 3 times a day for at least 7 full days..
- Acuvail** (Anti-inflammatory) – 1 drop in each operative eye 2 times a day as needed for pain for 48 hours.
- FML** (Steroid – shake well before using) – 1 drop in each operative eye:
 - 4 times a day for 1 month
 - 3 times a day for 1 month
 - 2 times a day for 2 weeks
 - 1 time a day for 2 weeks
- Refresh Tears** - Use 1 drop in the surgical eye(s) 4x/day for at least 2 months.

To avoid washing out the previously applied drops, **wait approximately 1 minute between drops.**
***If you receive generic eye drops, the bottles may look different than the pictures shown above.**